

8-02-00

A

08/01/00
1c867 U.S. PTO

1c867 U.S. PTO
09/629341
00/01/00

Please type a plus sign (+) inside this box → ☐

Approved for use through 09/30/2000. OMB 0651-0032
Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

UTILITY PATENT APPLICATION TRANSMITTAL <small>(Only for new nonprovisional applications under 37 C.F.R. § 1.53(b))</small>	Attorney Docket No.	204,728
	First Inventor or Application Identifier	Bong-Ho LEE
	Title	NOVEL MATERIAL SEPARATED FROM ECKLONTA CAVA, METHOD FOR EXTRACTING AND PURIFYING THE
	Express Mail Label No.	EK051386875US

APPLICATION ELEMENTS <small>See MPEP chapter 600 concerning utility patent application contents.</small>	ADDRESS TO: Assistant Commissioner for Patents Box Patent Application Washington, DC 20231	
1. <input checked="" type="checkbox"/> * Fee Transmittal Form (e.g., PTO/SB/17) (Submit an original and a duplicate for fee processing)	5. <input type="checkbox"/> Microfiche Computer Program (Appendix)	
2. <input checked="" type="checkbox"/> Specification [Total Pages 21] (preferred arrangement set forth below) <ul style="list-style-type: none">- Descriptive title of the invention- Cross References to Related Applications- Statement Regarding Fed sponsored R & D- Reference to Microfiche Appendix- Background of the invention- Brief Summary of the invention- Brief Description of the Drawings (if filed)- Detailed Description- Claim(s)- Abstract of the Disclosure	6. Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary) <ul style="list-style-type: none">a. <input type="checkbox"/> Computer Readable Copyb. <input type="checkbox"/> Paper Copy (identical to computer copy)c. <input type="checkbox"/> Statement verifying identity of above copies	
3. <input type="checkbox"/> Drawing(s) (35 U.S.C. 113) [Total Sheets]	ACCOMPANYING APPLICATION PARTS 7. <input checked="" type="checkbox"/> Assignment Papers (cover sheet & document(s)) 8. <input type="checkbox"/> 37 C.F.R. § 3.73(b) Statement (when there is an assignee) <input checked="" type="checkbox"/> Power of Attorney 9. <input type="checkbox"/> English Translation Document (if applicable) 10. <input type="checkbox"/> Information Disclosure Statement (IDS)/PTO-1449 <input type="checkbox"/> Copies of IDS Citations 11. <input type="checkbox"/> Preliminary Amendment 12. <input checked="" type="checkbox"/> Return Receipt Postcard (MPEP 503) (Should be specifically itemized) * Small Entity <input type="checkbox"/> Statement filed in prior application, Status still proper and desired 13. <input checked="" type="checkbox"/> Statement(s) (PTO/SB/09-12) 14. <input checked="" type="checkbox"/> Certified Copy of Priority Document(s) (if foreign priority is claimed) 15. <input type="checkbox"/> Other: _____	
4. Oath or Declaration [Total Pages 2] <ul style="list-style-type: none">a. <input checked="" type="checkbox"/> Newly executed (original or copy)b. <input type="checkbox"/> Copy from a prior application (37 C.F.R. § 1.63(d)) (for continuation/divisional with Box 16 completed)<ul style="list-style-type: none">i. <input type="checkbox"/> DELETION OF INVENTOR(S) Signed statement attached deleting inventor(s) named in the prior application, see 37 C.F.R. §§ 1.63(d)(2) and 1.33(b).		
* NOTE FOR ITEMS 1 & 13: IN ORDER TO BE ENTITLED TO PAY SMALL ENTITY FEES, A SMALL ENTITY STATEMENT IS REQUIRED (37 C.F.R. § 1.27), EXCEPT IF ONE FILED IN A PRIOR APPLICATION IS RELIED UPON (37 C.F.R. § 1.28).		
16. If a CONTINUING APPLICATION, check appropriate box, and supply the requisite information below and in a preliminary amendment: <input type="checkbox"/> Continuation <input type="checkbox"/> Divisional <input type="checkbox"/> Continuation-in-part (CIP) of prior application No: _____ Prior application information: Examiner _____ Group / Art Unit: _____ For CONTINUATION or DIVISIONAL APPS only: The entire disclosure of the prior application, from which an oath or declaration is supplied under Box 4b, is considered a part of the disclosure of the accompanying continuation or divisional application and is hereby incorporated by reference. The incorporation can only be relied upon when a portion has been inadvertently omitted from the submitted application parts.		

17. CORRESPONDENCE ADDRESS					
<input type="checkbox"/> Customer Number or Bar Code Label			or <input checked="" type="checkbox"/> Correspondence address below		
(Insert Customer No. or Attach bar code label here)					
Name	ADELMAN FRAYNE & SCHWAB Attorneys at Law				
Address	150 East 42nd Street New York, NY 10017				
City	State	Zip Code			
Country	Telephone	Fax			
U.S.A.	(212) 949-9022	(212) 949-9190			

Name (Print/Type)	Jay S. Cinamon	Registration No. (Attorney/Agent)	24,156
Signature	Jay S. Cinamon	Date	August 1, 2000

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.

*) SAME, AND USE THEREOF AS ANTIOXIDANTS

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICANT :

SERIAL NO. :

EXAMINER :

FILED :

GROUP NO. :

FOR (TITLE): NOVEL MATERIAL SEPARATED FORM ECKLONIA CAVA, METHOD FOR
EXTRACTING AND PURIFYING THE SAME, AND USE THEREOF AS
ANTIOXIDANTS

VERIFIED STATEMENT AS SMALL ENTITY

Hon. Commissioner of Patents and Trademarks
Washington, D.C. 20231

S I R:

The undersigned declare(s):

Exclusive rights in the above-identified inven
defined and named below, and "small entity" fees are apprc
entity is based upon the following:

☐

INDEPENDENT INVENTOR

An independent inventor is any inventor who:

- 1) has not assigned, granted, conveyed, or licensed, and
- 2) is under no obligation under contract or law to assign
rights in the invention to any person who could not likewise be classified as an independent
inventor if that person had made the invention, or to any concern which would not qualify as
a small business concern or a non-profit organization as defined in Rule 1.9.

☒

SMALL BUSINESS CONCERN

A small business concern is defined as a business concern:

- 1) whose number of employees, including those of its affiliates, does not exceed 500
persons, and
- 2) which has not assigned, granted, conveyed, or licensed, and is under no obligation under
contract or law to assign, grant, convey or license, any rights in the invention to any person who
could not be classified as an independent inventor if that person had made the invention, or to
any concern which would not qualify as a small business concern or a nonprofit organization as
defined in Rule 1.9. Concerns are affiliates of each other when, either directly or indirectly,
one concern controls or has the power to control the other, or a third party controls or has the
power to control both. The number of employees of the business concern is the average over
the fiscal year of the persons employed during each of the pay periods of the fiscal year.
Employees are those persons employed on a full-time, part-time or temporary basis during the
previous fiscal year of the concern.

STATEMENT OF FILING BY
EXPRESS MAIL (37 CFR § 1.10)
This correspondence is being
deposited with the United States
Postal Service on
August 1, 2000 in an
envelope as "Express Mail Post
Office to Addressee" Mailing
Label Number
EK05138687SUS addressed to
the Commissioner of Patents and
Trademarks, Washington, D.C. 20231

☐**NONPROFIT ORGANIZATION** (Check additional applicable box)

A nonprofit organization is defined as:

☐

1) a university or other institution of higher education located in any country; or

☐

2) a organization of the type described in section 501(c)(3) of the Internal Revenue Code of 1954 (26 U.S.C. 501(c)(3)) and exempt from taxation under Section 501(a) of the Internal Revenue Code (26 U.S.C. 501(a)); or

☐

3) any nonprofit scientific or educational organization qualified under a nonprofit organization statute of a state of the United States (35 U.S.C. 201(i)); or

☐

4) any nonprofit organization located in a foreign country which would qualify as a nonprofit organization under paragraphs (e)(2) or (3) of Rule 1.9 if it were located in the United States.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

VENTREE CO., LTD

(Print or type name of small entity)

* H. W. Lee(Signature and title of person authorized
to act on behalf of small entity)

, President

(Date)

July 28, 2000Seowoo-building 14, Yeoksam-dong 837-12,

(Address)

Gangnam-ku, Seoul, Republic of Korea

* If "small entity" is more than one inventor or a combination of "small entities", add name(s), address(es) and signature(s) of other parties below. Please include date.

Full Name of Second Joint Inventor, If Any	Inventor's Signature	Date
Residence	Citizenship	
Post Office Address		
Full Name of Third Joint Inventor, If Any	Inventor's Signature	Date
Residence	Citizenship	
Post Office Address		
Full Name of Fourth Joint Inventor, If Any	Inventor's Signature	Date
Residence	Citizenship	
Post Office Address		

(page 2 of 2)

NOVEL MATERIAL SEPARATED FROM *Ecklonia cava*, METHOD FOR
EXTRACTING AND PURIFYING THE SAME, AND USE THEREOF AS
ANTIOXIDANTS

5

BACKGROUND OF THE INVENTION

1. Field of the Invention

10 This invention relates to materials separated from
Ecklonia cava, their preparation, and their use as
antioxidants. More particularly, the present invention
relates to materials extracted from *Ecklonia cava*, which is
superior in antioxidative activity and thermal stability, a
method for the extraction and purification of such
15 antioxidative materials from *Ecklonia cava*, and use thereof
as antioxidants.

2. Description of the Prior Arts

20 The human body can be maintained alive, utilizing the
energy obtained from nutrients through aerobic metabolisms.
However, various physical, chemical and biological stresses
change oxygen, used as an electron acceptor in aerobic
metabolisms, into harmful active oxygen species such as
25 superoxide anion radical (O_2^-), hydrogen peroxide, or

hydroxy radical to generate fatal physiological disorders or to induce diseases in the human body. The human body possesses an antioxidative mechanism as a self-defense mechanism to scavenge such active oxygen species. However, 5 active oxygen species, if they occur with more potential power than the defense ability of the body, break the factors responsible for the immune system, such as proteins, DNA, enzymes and T cells, to generate disorders. Also, such powerful active oxygen species attack unsaturated fatty 10 acids, which are constituents of cellular membranes, to cause a peroxidation reaction. It is known that lipid peroxides accumulated in the body cause aging and disorders.

There has been widely acknowledged the theory that aging and adult diseases are attributed to active oxygen 15 species. Researches have been directed to the theory since the report on auto-oxidation in the 1940s. Antioxidants, able to inhibit an oxidation reaction, can be used to inhibit acidification of foods, aging of the human body and the like. In particular, synthetic antioxidants, such as 20 butylhydroxytoluene (BHT) and butylhydroxyanisole (BHA), have been extensively used in the food industry. However, these synthetic materials are known to be carcinogenic, so that humans are reluctant to eat such synthetic antioxidant-containing foods. An intensive and extensive interest has 25 been taken in methods for extracting antioxidative materials

from natural materials because the extracts do not cause cancers. Of them, a method for extracting beta-carotene from carrots is found to be economically unfavorable because its output is very small. Recently, Shin-Ya et al. have
5 extracted and identify naphtherpin, a kind of antioxidant, from Streptomyces CL 190 (K. Shin-Ya, S. Imai, K. Furihata, Y. Hayakawa, K. Kato, G. D. Vanduyne, J. Clardy and H. Seto, J. Antibiotics 43, 444(1990)), which, however, was not commercialized. Further, Teshima et al. and Mo et al.
10 reported the extraction of antioxidative materials from microorganisms (Y. Techima and K. Shin-Ya, J. Antibiotics 44, 685(1991); C. J. Mo, K. Shin-Ya, K. Furihata, A Shimazu, Y. Hayakawa and H. Seto, J. Antibiotics 43, 1337(1990)), but it was not commercialized, either. In addition, active
15 research efforts have been and continue to be directed to methods for extracting various antioxidative materials from farm products and marine products. Particularly, tocopherol is well-known as an antioxidant, and tea extracts are known to contain various antioxidants. Korean Pat. Publication No.
20 1997-3067 refers to a method for preparing natural antioxidants by immersing fish skin in hot water to extract gelatin and hydrolyzing the extracted gelatin in a three-step enzyme membrane reactor to obtain enzymatic lysates.

Also, Korean Pat. Application No.99-60007 refers to a
25 method for preparing a natural antioxidant from wild roses,

in which the flowers are immersed in an organic solvent to obtain an extract of antioxidative activity from which beta-glucogalin is isolated and purified.

Meanwhile, other attempts to extract physiologically
5 active materials, especially antioxidative materials from seaweeds have been made since late 1980s and have met with success in France and Japan. Tagaki and Miyashida reported that natural compounds extracted from 12 kinds of seaweeds in Japan waters contain tocopherol consisting mainly of an
10 alpha-type along with a minor portion of a beta-type (Miyashita and T.Tagaki, Agric. Biol. Chem. 51, 3115(1987)). Also, Kaneniwa et al. successfully extracted lipid materials of antioxidative activity from seaweeds and identified them as 5-olefinic acids, which are unusual materials in regard
15 to antioxidants (M. Kaneniwa, Y. Itabashi and T Tagaki, Nippon Suisan Gakkashi 53, 861(1987)). Nishibori and Nakami reported that antioxidative lipid materials were extracted from seven kinds of seaweeds with a hexane/ethanol mixture and the lipid materials extracted from lavers and brown
20 seaweeds have an antioxidative activity similar to that of BHA and alpha-tocopherol (S. Nishibori and K. Namiki, kateigaku zaxtusi 36, 17(1985)). But the extracted amount was so small that this method was not commercialized.

It is reported that methanol and chloroform extracts
25 from lavers and brown seaweeds and sea tangles are superior

in antioxidative activity to BHA (Jae-Han Park, Gyu-chan Kang, Sang-Bong Paek, Yun-Hyung Lee, Gyu-Soon Lee, Korean Journal of Food Science and Technology 23, 256(1991)). They used methanol and chloroform, in order, for the extraction
5 of the antioxidative materials from 12 kinds of seaweeds, but failed to commercialize the materials owing to their weak thermal stability.

SUMMARY OF THE INVENTION

10


In view of these situations, the present inventors have made an extensive research designed to extract and purify useful materials from seaweeds. As a result, the present invention has been completed through the development
15 of novel materials separated from *Ecklonia cava*, which can be used as antioxidants because they have both an excellent antioxidative activity and a thermal stability. In connection, there was also developed a method for extracting and purifying the novel materials of antioxidative activity.

20 Accordingly, it is an object of the present invention to provide novel materials extracted from *Ecklonia cava*, a kind of seaweed in Korean waters.

It is another object of the invention to provide a method for extracting and purifying such natural materials
25 from *Ecklonia cava*.

It is a further object of the invention to provide use of such natural materials as antioxidants by taking advantage of their superior radical scavenging activity and thermal stability.

5 Novel materials for achieving said object are represented by the following Formula I.



I

wherein R is hydrogen or a hydroxy group.

10 To accomplish another object of the present invention, the method for extracting and purifying said materials from *Ecklonia cava*, comprises the following steps; extracting antioxidative ingredients from powdered *Ecklonia cava* one or more times with an organic solvent; fractionating the
15 antioxidative ingredients one or more times in solvents; and purifying the solvent fractions by chromatography.

To accomplish another object of the present invention,


novel materials extracted from *Ecklonia cava* are used as antioxidants.

DETAILED DESCRIPTION OF THE INVENTION

5

Novel materials according to the present invention are represented by the following Formula I, and can be used as antioxidants by virtue of excellent scavenging activity and thermal stability.

10



I

wherein R is hydrogen or a hydroxy group.

The antioxidative materials is extracted and purified
15 as follows.

Ecklonia cava is washed with distilled water to eliminate impurities, dried in the absence of direct

sunlight, and crushed into powder. At room temperature, this powdered *Ecklonia cava* is immersed in an organic solvent to obtain an extract containing the compound of Formula I. Useful in this extraction is an organic solvent
5 which is preferably selected from the group consisting of methanol, ethanol, ethyl acetate, acetonitrile, acetone, water and a mixture thereof and water/ethanol mixture. Optionally, said extraction procedure may be conducted at least twice in order to increase the yield, wherein a
10 different organic solvent may be used in each repetition step. For eliminating undesired materials and the solvent from the extracts, use can be made of separation and concentration instruments such as a centrifuge and a rotary evaporation concentrator. The extracts obtained through
15 said procedure can be directly used in various fields of applications, so that they are economically favorable because of no additional processes. In the case that the material has to be of high purity, it is preferred that the following additional separation and purification steps are
20 conducted.

The materials obtained from the extraction step are treated in the next fractionation step. Optionally, the method may further comprise the step of dissolving the extracts in ethyl acetate and/or methanol and removing
25 undissolved residues before the fractionation step. By way

of examples, and not limitation, the solvent-fractionating step can be conducted in a three-step manner. For a primary fractionation, an aqueous, 10 to 90 % methanol solution is used as a polar layer while, as a nonpolar layer, a linear or cyclic hydrocarbon solvent, such as hexane, cyclohexane or pentane and an aromatic solvent, such as benzene or toluene, are used alone or in mixed combinations thereof. In particular, it is preferred to use an aqueous 60 to 90 % methanol solution and hexane. As a result of the primary solvent-fractionation, the novel compound dominantly exists in the aqueous methanol solution layer. In a secondary solvent-fractionation, the aqueous solution containing the novel compound is subjected to solvent extraction using an aqueous 10 to 60 % methanol solution as a polar layer and one or more ethers including isopropylether as a nonpolar layer. An aqueous 20 to 40 % methanol solution and isopropylether are preferred. Likewise, most of the novel compound is found to exist in the aqueous 10 to 60 % methanol solution layer. For a tertiary solvent-fractionation, an aqueous 10 to 60 % methanol solution is used as a polar layer while, as a nonpolar layer, chloroform and dichloromethane are used alone or in mixed combinations thereof. Preferably, an aqueous 30 to 50 % methanol solution and chloroform are used.

For purifying the organic fraction obtained through

the above procedure, the aqueous methanol solution layer is re-dissolved in pure distilled water and then passed through membranes to separate active ingredients. Of them, highly active ingredients are collected and purified by, for example, medium pressure liquid chromatography (MPLC) or high performance liquid chromatography (HPLC). In regard to the purification, other chromatographic techniques may be used if necessary.

The antioxidative activity of the novel compound obtained through the extraction and purification process is evaluated by measuring its radical scavenging activity against 1,1-diphenyl-2-picrylhydrazyl (DPPH), which has free radicals attached thereto, according to the Blois method. A thermal stability of the compound of interest can be evaluated by measuring its antioxidative activity at plural temperature points.

A better understanding of the present invention may be obtained in light of the following examples which are set forth to illustrate, but are not to be construed to limit the present invention.

EXAMPLE 1: Extraction Step

410 g of *Ecklonia cava*, which was dried out of direct sunlight and then crushed into powder, was dissolved in 6.5L

of methanol in a 1 L round bottom flask and the solution was slowly stirred for 12 hours at room temperature to extract antioxidative materials. Thereafter, the methanol extract was centrifuged at a low temperature to remove undesired
5 impurities, followed by the removal of the methanol with the aid of a rotary evaporation concentrator. The concentrated extract was measured to be 64.1 g.

EXAMPLE 2: Fractionation Step

10

The extract obtained in Example 1 was dissolved in 4 L of ethyl acetate and concentrated by filtration and then, the undissolved residue was removed. The ethyl acetate solution was subjected to the first solvent-fractionation
15 step of using 1 L of an aqueous 90 % methanol solution and 3.5 L of n-hexane. As a result of this fractionation, antioxidative active ingredients were found in the aqueous methanol solution layer. Again, this aqueous solution layer was subjected to the second solvent-fractionation step of
20 using 1 L of an aqueous 30 % methanol solution and 1 L of isopropylether. Likewise, the antioxidative active ingredients were in the aqueous 30 % methanol solution layer. The third solvent-fractionation step was performed with the aqueous 30 % methanol solution layer using 1 L of an aqueous
25 40 % methanol solution and 1 L of chloroform. Thereafter,

the aqueous 40 % methanol solution layer was dried in vacuo, to give 2.85 g of an organic fraction.

EXAMPLE 3: Purification Step

5

The organic fraction obtained in Example 2 was loaded in a 25x500 mm glass column filled with ODS (octadecylsilyl) resin with a diameter of 200 μm and was eluted with a 30 % methanol solution to give 850 mg of active ingredients. 10 These active ingredients were subjected to high performance liquid chromatography (acetonitrile:water = 20:80, flow rate=2.0 ml/min, 10x250 mm C-18 column) to give a pure material. Various spectrophotometric analyses were carried out. The results are as follows:

15

Ultraviolet-visible spectra: UV (MeOH) λ_{max} 231nm (ϵ 6300), 246, 295 (8800);

Infrared spectra: IR (film) ν_{max} 3300 (OH), 2950, 1590 (aromatic) cm^{-1} ;


20

Mass spectra: HRFABMS (pos) m/z 745.1039 [$(M+H)^+$, $\text{C}_{36}\text{H}_{24}\text{O}_{18}$, $\Delta+0.3\text{mmu}$]; and

^1H -NMR: δ ^1H (multi, JHz) 6.33(H, d, 1.6), 6.61(H, d, 1.6), 6.64(H, s), 6.64(H, s), 6.64(H, s), 6.75(H, s), 6.80(H, s), 6.79(H, s), 6.79(H, s), 6.22(H, d, 1.2), 6.66(H, d, 1.2).

25 From these data, the structure of the material was

identified as the following Formula Ia (Dicaval A).



Ia

The other was also analyzed in various spectroscopy.


5 The results are as follows:

Ultraviolet-visible light spectra UV (MeOH) λ_{\max} 232nm:
(ϵ 6500), 246, 296 (9000);

Infrared light spectra: IR (film) ν_{\max} 3350 (OH), 2950,
1580 (aromatic) cm^{-1} ;

10 Mass spectra HRFABMS (pos) m/z : 761.1024 [(M+H)⁺,
C₃₆H₂₄O₁₉, Δ +0.4mmu] ;

¹H-NMR δ ¹H(multi, JHz): 6.33(H, d, 1.6), 6.61(H, d,
1.6), 6.64(H, s), 6.64(H, s), 6.64(H, s), 6.75(H, s), 6.79(H,
s), 6.79(H, s), 6.22(H, d, 1.2), 6.66(H, d, 1.2). From
15 these data, the structure of the material was identified as
the following Formula Ib (Dicaval B).



Ib

EXAMPLE 4: Measurement of Antioxidative Activity

5 Dicaval A and Dicaval B obtained in Example 3 were measured for antioxidative activity according to the Blois method. For this, an examination was made of the radical scavenging activity of the two compounds against 1,1-diphenyl-2-picrylhydrazyl (DPPH) to which free radicals were

10 attached. First, 20 mg of DPPH was dissolved in 150 ml of ethanol to prepare a DPPH solution. To 600 μ l of the DPPH solution was added with 250 μ l of dimethylsulfoxide (DMSO), diluted with an appropriate amount of ethanol and shaken for

15 10 seconds, after which this control was adjusted, in absorbance at 517 nm, from 0.94 to 0.97. To 1 ml of the DPPH solution which was adjusted from 0.94 to 0.97 likewise,

each of the samples (μg to mg) was added and then, reacted for 10 min, followed by measuring the absorbance of the solution. The antioxidative activity of each of the samples was determined by the DPPH radical scavenging activity which was represented by a reduced absorbance compared with that of the control.

The antioxidative activity was compared between the conventional antioxidants and the compounds of the invention and the results are given in Table 1, below.

TABLE 1

Sample amount (μg)	BHT	Dicaval A	Dicaval B	Ascorbic acid
10	74 %	100 %	100 %	100 %
20	87 %	100 %	100 %	100 %
100	91 %	100 %	100 %	100 %

EXAMPLE 5: Evaluation of Thermal Stability

50 μg of each of Dicaval A and Dicaval B obtained in Example 3 was heated at 40°C, 60°C, 80°C, and 100°C for 1 hour to measure antioxidative activity. The results are shown in Table 2, below.

TABLE 2

Antioxidativity (Absorb. change)

Temp. (°C)	Dicaval A	Dicaval B	Notes
40	0.87	0.88	Stable
60	0.89	0.88	Stable
80	0.88	0.87	Stable
100	0.90	0.89	Stable


From the results of the above tables, it can be said that the antioxidative natural materials of the present invention are superior in antioxidative activity to BHT and excellent, in particular, in thermal stability in the aspect that their antioxidative effects are constantly maintained in a broad range of temperatures.

Over conventional antioxidants, the novel materials of the present invention have advantages of being superior in scavenging activity and thermal stability and showing minimal side effects when administered. Thus, the novel materials can replace the conventional antioxidants.

Whereas particular embodiments of this invention have been described above for purposes of illustration, it will be evident to those skilled in the art that numerous variations of the details of the present invention may be made without departing from the invention as defined in the appended claims.

What is claimed is:

1. A material represented by the following Formula I.




I

wherein R is hydrogen or a hydroxy group.

2. An extract containing the material of claim 1, obtained from *Ecklonia cava* by use of an organic solvent.

3. A method for extracting and purifying from *Ecklonia cava*, the materials, represented by the following Formula I:



I

wherein R is hydrogen or a hydroxy group,
comprising the steps of:

- 5 extracting antioxidative ingredients from *Ecklonia*
cava once or more times with an organic solvent;
 fractionating the antioxidative ingredients one or
more times in solvents; and
 purifying the solvent fractions by chromatography.

10

4. The method as set forth in claim 3, wherein said
organic solvent is selected from the group consisting of
methanol, ethanol, ethyl acetate, acetonitrile, acetone,
aqueous solutions thereof, and mixtures thereof.

15

5. The method as set forth in claim 3, wherein the
extracting step is repeated using the same or different

solvents.

6. The method as set forth in claim 3, wherein the fractionating step comprises:

- 5 a primary solvent-fractionating step of fractionating the extract by using an aqueous 10 to 90 % methanol solution as a polar layer, and a linear or cyclic hydrocarbon solvent, an aromatic solvent, or a mixture thereof as a nonpolar layer;
- 10 a secondary solvent-fractionating step of fractionating an aqueous methanol layer obtained in the primary step by using an aqueous 10 to 60 % methanol solution as a polar layer and at least one ether as a nonpolar layer; and
- 15 a tertiary solvent-fractionating step of fractionating an aqueous methanol layer obtained in the secondary step by using an aqueous 10 to 60 % methanol solution as a polar layer and chloroform, dichloromethane, or a mixture thereof as a nonpolar layer.

20

7. The method as set forth in claim 3, wherein the fractionating step comprises:

- a primary solvent-fractionating step of fractionating the extract by using an aqueous 10 to 90 % methanol
- 25 solution as a polar layer, and hexane as a nonpolar

layer;

5 a secondary solvent-fractionating step of
fractionating an aqueous methanol layer obtained in
the primary step by using an aqueous 20 to 40 %
methanol solution as a polar layer and isopropyl
ether as a nonpolar layer; and

10 a tertiary solvent-fractionating step of fractionating
an aqueous methanol layer obtained in the secondary
step by using an aqueous 30 to 50 % methanol solution
as a polar layer and chloroform as a nonpolar layer.

8. The method as set forth in claim 3, further
comprising the step of dissolving the extract in ethyl
acetate and/or methanol and providing the dissolved portion
15 to the fractionating step.

9. The method as set forth in claim 3, wherein the
chromatography is medium pressure liquid chromatography
(MPLC) or high performance liquid chromatography (HPLC).
20


10. Use of the material of claim 1 as antioxidants.

11. Use of the extract of claim 2 as antioxidants.

ABSTRACT

Disclosed are novel materials separated from *Ecklonia* cava, a method for extracting and purifying the same, and
5 the use thereof for antioxidants. The method comprises extracting antioxidative ingredients from powdered *Ecklonia* cava one or more times with an organic solvent; fractionating the antioxidative ingredients one or more
10 times in solvents; and purifying the solvent fractions by chromatography. Superior in scavenging activity and thermal stability, the extract from *Ecklonia cava* can be used as antioxidants and is suitable in commercialization.

United States Patent & Trademark Office
Office of Initial Patent Examination -- Scanning Division



Application deficiencies were found during scanning:

☐ Page(s) _____ of Declaration were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ Scanned copy is best available.